arXiv:1404.4010v2 [hep-th] 9 Jan 2015

Duality between zeroes and poles in holographic systems with massless fermions and a dipole coupling

James Alsup,[1,*] Eleftherios Papantonopoulos,[2,†] George Siopsis,[3,‡] and Kubra Yeter[3,§]

[1] *Computer Science, Engineering and Physics Department, The University of Michigan-Flint, Flint, MI 48502-1907, USA*

[2] *Department of Physics, National Technical University of Athens, Zografou Campus GR 157 73, Athens, Greece*

[3] *Department of Physics and Astronomy, The University of Tennessee, Knoxville, TN 37996 - 1200, USA*

(Dated: January 12, 2015)

Abstract

We discuss the zeroes and poles of the determinant of the retarded Green function ($\det G_R$) at zero frequency in a holographic system of charged massless fermions interacting via a dipole coupling. For large negative values of the dipole coupling constant p, $\det G_R$ possesses only poles pointing to a Fermi liquid phase. We show that a duality exists relating systems of opposite p. This maps poles of $\det G_R$ at large negative p to zeroes of $\det G_R$ at large positive p, indicating that the latter corresponds to a Mott insulator phase. This duality suggests that the properties of a Mott insulator can be studied by mapping the system to a Fermi liquid. Finally, for small values of p, $\det G_R$ contains both poles and zeroes (pseudo-gap phase).

PACS numbers: 11.15.Ex, 11.25.Tq, 74.20.-z

*Electronic address: jalsup@umflint.edu

†Electronic address: lpapa@central.ntua.gr

‡Electronic address: siopsis@tennessee.edu

§Electronic address: kyeter@tennessee.edu

I. INTRODUCTION

The AdS/CFT correspondence is a principle that connects a strongly coupled d-dimensional conformal field theory with a weakly coupled $(d+1)$-dimensional gravity theory [1]. This principle, also known as the gauge/gravity duality, is well-founded in string theory, and has been applied to many field theories having gravity duals with the most noticeable application in condensed matter physics. One of the earliest applications of the gauge/gravity duality to condensed matter systems was the holographic description of a superconductor [2]. It is described by an Einstein-Maxwell-scalar field theory in which the breaking of a gauge symmetry in the gravity sector corresponds, in the boundary theory, to an operator (the order parameter) which condenses below a critical temperature signaling the onset of superconductivity.

The gauge/gravity duality has also been used as a framework for Fermi liquid behavior [3], non-linear hydrodynamics [4], quantum phase transitions [5] and transport [6]. Thus, classical gravity theories have been transformed into a laboratory for exploring condensed matter physics from a different point of view [7]. In this way the holographic principle is a powerful tool for studying strongly-correlated systems.

Recently, there has been considerable activity using the holographic principle to describe metallic states of matter outside of Landau's Fermi liquid theory [8, 9]. Experiments provide evidence for the existence of materials whose electronic structure cannot be described by Fermi liquid theory, nor by any other known effective theory. While Fermi liquids are characterized by stable quasi-particles, these exotic materials possess well-defined Fermi surfaces but lack long-lived quasi-particles. A quantitative discussion of signatures of different excitations is presented in detail below. To describe such metallic systems, one needs to introduce charged fermions in the gravity sector of the holographic system. However, in general, fermions cannot be treated classically as in the case of charged scalar fields of holographic superfluids or superconductors. The coupled system of Einstein-Maxwell-Dirac field equations become tractable only in the limit in which the fermions may be treated locally in the bulk as a charged ideal fluid of zero temperature. The solutions of these systems are known as an electron star [10–12], Dirac hair [13] or confined Fermi liquid [14].

In another approach to describe the various phases of a metallic state at low temperatures, a dipole coupling for massless charged fermions is introduced [15, 16]. The modified Dirac

equation in the background of an AdS-Reissner-Nordström black hole produces a boundary fermionic spectrum with vanishing spectral weight for a range of energies around $\omega = 0$ without the breaking of any symmetry. As the dipole coupling strength is varied, the fluid has Fermi, marginal Fermi, non-Fermi liquid phases and also an additional insulating phase. The insulating phase possesses the characteristics representative of Mott insulators, namely the dynamic formation of a gap and spectral weight transfer. Similar behavior was found in other constructions [17–21]. The dependence of the Fermi surface and selection of (non-) Fermi liquid type are discussed in [22].

For a Fermi liquid, the characteristic signature of a quasi-particle is a pole in the retarded Green Function G_R. The location of the pole

$$G_R^{-1}(\omega = 0, k = k_F) = 0 \ , \tag{1}$$

in momentum space at zero frequency dictates the Fermi surface and dispersion. Additionally, the real component of the Green function, $\Re G_R$, must change signs. In a normal, weakly-coupled metal, this is only possible through the pole, and via Luttinger's theorem, integrating over the positive region of G_R one obtains the particle density [23].

Apart from Fermi liquids, one of the principal examples of a strongly correlated condensed matter system is the Mott insulator, parent state of high-temperature copper-oxide superconductors in which a gap is dynamically generated. An explanation for the insulating nature [24] is that a Mott gap manifests as a surface of zeroes in the Green function. The Green function of interest, G_R, may be expressed in terms of the non-interacting theory Green function, G_0, and the self-energy Σ,

$$G_R^{-1}(\omega, k) = G_0^{-1}(\omega, k) + \Sigma(\omega, k) \ , \tag{2}$$

of which a zero is indicative of a divergent self-energy. It is the divergent self-energy that restricts the band from crossing the Fermi energy leading to Mottness. The full condition for Mottness is specifically on the eigenvalues of the matrix G_R and represented with the determinant as,

$$\det G_R(\omega = 0, k = k_L) = 0 \ . \tag{3}$$

For our purposes, we are interested in the spinor space but the statement may be extended to multiple bands [25]. A mean-field theory approach such as conventional BCS theory as

calculated with the Nambu propagator, is unable to produce such a result. The location of poles and zeroes in G_R provides a path to uncover a system's behavior.

The interplay between poles and zeros is a characteristic feature of QCD. The renormalization from high (UV) to low (IR) energy of couplings and masses in QCD involves free quarks at UV scales, which exhibit poles in the propagator, while at IR scales the poles are converted to zeros [26] signaling the formation of bound states. The zero of the single quark propagator in the IR is due to an orthogonal configuration between the (UV) single quark description and the composite mesons in the IR.

In condensed matter physics the competition of poles ($k = k_F$) and zeroes ($k = k_L$) in the Green function provides an illuminating criterion for characterizing the different phases found in strongly-coupled fermion systems [25]. Exploring the Hubbard model in this context has led to a variety of features present in doped Mott insulators, including spectral weight transfer, Fermi arcs found in the pseudo-gap state, and Fermi pockets [24, 27]. The Fermi arcs may be seen as a general consequence of a composite state of UV fields, and hence zeroes, existing at low energies within the Hubbard model [28]. In [29], a "pole-zero mechanism" for the two-dimensional Hubbard model was used to characterize the transition from metal to Mott insulator in terms of interfering pole and zero surfaces. The formalism is comprehensive enough to accommodate arcs, hole pockets, in-gap states, Lifshitz transitions and non-Fermi liquids. The work was furthered in [30] to address several anomalies in connection to (non) d-wave symmetries seen in the pseudo-gap and to suggest a topological quantum phase transition responsible for a non-Fermi liquid phase.

A holographic description of the structure of poles and zeros and their interconnection addresses central issues in strongly-coupled fermionic systems. A Fermi surface in the Green function for holographic fermions was discovered in [8, 32] with the discussion further expanded in [3, 9]. Near the Fermi surface, the structure of the poles is expanded similarly to eq. (2) as

$$G_R^{-1}(\omega, k) \sim k - k_F - \omega/v_F - \Sigma(\omega, k) \tag{4}$$

with self-energy $\Sigma(\omega, k)$ [33] given as

$$\Sigma(\omega, k) \sim c\omega^{2\nu_k^{\pm}} \quad , \quad 6\nu_k^2 = k^2 - \frac{q^2}{2} \ , \tag{5}$$

with fermion charge q and numerically calculated constant c. It is this scaling dimension ν_k that selects the leading term in (4) dictating the dispersion relation at the Fermi surface, and

hence the type of fluid. For scaling dimension $\nu_k < 1/2$, the pole of G_R corresponds to an unstable quasi-particle. This is identified as a non-Fermi fluid. With the value $\nu_k = 1/2$, the excitations are of marginal Fermi fluid type. For $\nu_k > 1/2$ the dispersion relation is linear. This is the Fermi (non-Landau) fluid. Lastly, imaginary ν_k corresponds to "log oscillatory" solutions.

Since the discovery of Fermi surfaces within the strongly coupled field theories analyzed via holography, an exploration of fermions has taken place. Mechanisms underlying different p-wave [34] and d-wave [35] symmetries for spectral functions have been offered. An effective lattice was added in [36] by using a modulated chemical potential. The spatial dependence alters the energy scaling and offers identifying signs for a holographic realization of a pseudo-gap state.

More fundamental (top-down) holographic theories from truncations of string/M-theory have produced viable models describing superfluity and superconductivity [37–41]. In higher dimensions, fermions are typically coupled to gravity and gauge fields with high-order derivatives. There are consistent truncations to lower dimensions in the fermionic sector giving interesting fermionic couplings [42, 43]. Dipole (Pauli) couplings are found to be a common feature of the theories with the coupling constant realized as scalar-dependent [44]. It would be interesting to explore their effect on fermion spectral functions at the boundary.

In this work, we aim at studying the zeroes and poles of the Green function G_R in a holographic system with a gravity sector consisting of an AdS-Reissner-Nordström black hole and a dipole coupling of massless charged fermions to an electromagnetic field. By exploring a duality between the zeroes and poles of this holographic system we show that by varying the dipole coupling strength, G_R may possess only poles, only zeroes, or both poles and zeroes.

This behavior suggests that the dipole coupling strength plays the role of an order parameter in the Mott physics. By studying a simple model in which a fermion is coupled to a gauge field through a dipole interaction in the bulk, we find that on the field theory side as the strength of this interaction is varied, a new band in the density of states emerges and spectral density is transferred between bands. In the language of Green functions only poles is the signature of the Fermi and non-Fermi liquids, only zeroes corresponds to the Mott insulating phase, and the coexistence of both poles and zeroes is the pseudo-gap phase.

This duality between zeroes and poles also suggests that within the holographic system,

the properties of the Green function G_R signifying the Mott phase can be inferred by mapping to the well understood Green function of a Fermi liquid by simply changing the sign of the dipole coupling constant $p \to -p$.

The discussion is organized as follows. In Section (II) we set up the theory and derive the Einstein-Maxwell-Dirac equations. In Section (III) we solve the Dirac equation. In Section (IV) we discuss the duality of poles and zeros, and finally, Section (V) contains our conclusions.

II. GRAVITATIONAL BULK

The bulk dynamics in an asymptotically Anti-de Sitter (AdS) space is described by the Einstein-Maxwell action with cosmological constant $\Lambda = -3/L^2$,

$$S = \int d^4x\sqrt{-g}\left[\frac{R+6/L^2}{16\pi G} - \frac{1}{4}F_{MN}F^{MN}\right] \ , \tag{6}$$

where $F_{MN} = \partial_M A_N - \partial_N A_M$ is the field strength of the $U(1)$ vector potential A_M. For convenience, we set $L = 4\pi G = 1$.

The Einstein-Maxwell equations admit a charged four-dimensional AdS black hole solution,

$$ds^2 = \frac{1}{z^2}\left[-h(z)dt^2 + \frac{dz^2}{h(z)} + dx^2 + dy^2\right] \ . \tag{7}$$

The metric function is given by

$$h(z) = 1 - \left(1+\mu^2\right)z^3 + \mu^2 z^4 \ , \tag{8}$$

with the horizon radius set at $z = 1$, and the $U(1)$ potential is

$$A_t = \mu\,(1-z) \ , \quad A_z = A_x = A_y = 0 \ , \tag{9}$$

corresponding to a non-vanishing electric field in the radial z direction,

$$F_{tz} = -F_{zt} = \mu \ . \tag{10}$$

The Hawking temperature is given by

$$T = -\frac{h'(1)}{4\pi} = \frac{3-\mu^2}{4\pi} \ , \tag{11}$$

with $\mu^2 = 3$ providing the zero temperature limit. There is a scaling symmetry of the solutions found as

$$z \ \to \lambda z \ , \quad x \to \lambda x \ , \quad \mu \ \to \mu/\lambda \ , \quad T \to T/\lambda \ , \tag{12}$$

and we should only report on scale-invariant quantities, such as T/μ, etc.

In this gravitational background we add a massless fermion with charge q and we include a dipole coupling to the $U(1)$ field [15, 16]. The action is

$$S_{\rm fermion} = i \int d^4x \sqrt{-g} \bar{\Psi} \left[\not{D} - p \Sigma^{MN} F_{MN} \right] \Psi \ . \tag{13}$$

The various terms in the action are

$$\begin{aligned} \not{D} &= e_a^M \Gamma^a \left(\partial_M + \Omega_M - iq A_M \right) \ , \\ \Omega_M &= \frac{1}{8} \omega_{abM} \left[\Gamma^a, \Gamma^b \right] \ , \\ \omega_{abM} &= \eta_{ac} \omega^c_{bM} \ , \\ \omega^a_{bI} &= e^a_M \partial_I e^M_b + e^a_M e^N_b \Gamma^M_{NI} \ , \\ \Sigma^{MN} &= \frac{i}{4} \left[\Gamma^a, \Gamma^b \right] e^M_a e^N_b \ , \end{aligned} \tag{14}$$

with spin connection ω_{abM} and vierbein e^M_a, and lower-case indices a, b belong to the tangent space.

In the conventional case, $p = 0$, the system is of two non-interacting Weyl fermions of opposite chiralities. With $p \neq 0$, the dipole term introduces an interaction between the two Weyl fermions. The system corresponds to an order parameter in the dual gauge theory of conformal dimension $\Delta = \frac{3}{2}$.

To analyze the Dirac equations of the bulk fermion we find it more convenient to go to momentum space by Fourier transforming

$$\Psi = e^{-i\omega t - ikx} \sqrt{\frac{z^3}{h}} \begin{pmatrix} \psi_- \\ \psi_+ \end{pmatrix} \quad , \quad \psi_\pm = \begin{pmatrix} \psi_{\pm 1} \\ \psi_{\pm 2} \end{pmatrix} \ , \tag{15}$$

and choosing a basis for the Γ matrices

$$\begin{aligned} \Gamma^1 &= \begin{pmatrix} i\sigma_1 & 0 \\ 0 & i\sigma_1 \end{pmatrix} \quad , \quad \Gamma^2 = \begin{pmatrix} -\sigma_2 & 0 \\ 0 & -\sigma_2 \end{pmatrix} \ , \\ \Gamma^3 &= \begin{pmatrix} 0 & -i\sigma_2 \\ i\sigma_2 & 0 \end{pmatrix} \quad , \quad \Gamma^4 = \begin{pmatrix} -\sigma_3 & 0 \\ 0 & -\sigma_3 \end{pmatrix} \ . \end{aligned} \tag{16}$$

The Dirac equation decomposes into decoupled equations

$$\begin{aligned} \pm h\psi'_{-,12} + \left(\mu q(1-z) + \omega \pm k\sqrt{h} \mp p\mu\sqrt{h}\right)\psi_{-,21} &= 0, \\ \pm h\psi'_{+,12} + \left(\mu q(1-z) + \omega \mp k\sqrt{h} \mp p\mu\sqrt{h}\right)\psi_{+,21} &= 0 \ . \end{aligned}$$

The presence of the dipole coupling p modifies the Dirac equation. The effects of the coupling will be seen more clearly in the solutions of the Dirac equations.

III. SOLUTIONS OF THE DIRAC EQUATIONS

To solve the Dirac equations we choose in-going boundary conditions at the horizon,

$$\psi_{\pm,12} = (1-z)^{-i\omega/(4\pi T)}\,\mathcal{F}_{\pm,12} \ . \tag{17}$$

The ratios

$$\xi_\pm = \frac{\mathcal{F}_{\pm 1}}{\mathcal{F}_{\pm 2}} \ , \tag{18}$$

satisfy the non-linear flow equations [9]

$$h\xi'_\pm + \left[\omega + \mu q(1-z) + \sqrt{h}\,(\pm k + \mu p z)\right]\xi_\pm^2 + \omega + \mu q(1-z) + \sqrt{h}\,(\mp k - \mu p z) = 0 \ , \tag{19}$$

together with the in-going boundary conditions,

$$\xi_\pm = \begin{cases} i & , \ \omega \neq 0 \ , \\ \\ i\sqrt{\frac{q/\sqrt{2}+\sqrt{3}p\pm k}{q/\sqrt{2}-\sqrt{3}p\mp k}} & , \ \omega = 0 \ . \end{cases} \tag{20}$$

The solution to the flow equations determines the retarded Green function as

$$G_R(\omega,k) = \begin{pmatrix} G_+ & 0 \\ 0 & G_- \end{pmatrix} \ , \quad G_\pm(\omega,k) = \xi_\pm|_{z\to 0} \ . \tag{21}$$

From the symmetries of the Dirac equation (15), we deduce the relation between the Green functions

$$G_\pm(\omega,k) = G_\mp(\omega,-k) \ . \tag{22}$$

The Fermi momentum is found as the pole

$$G_\pm^{-1}(\omega=0, k=k_F) = 0 \ . \tag{23}$$



FIG. 1: Scaling dimension ν_{k_F} *vs.* p for $q = 1$.

Since both solutions $\psi_\pm$ are normalizable, there is an alternative quantization in which G_+ and G_- are interchanged.[1] This leads to physically equivalent results.

To identify the type of (non-) Fermi fluid, we look at the near-horizon region, $z \to 1$. Employing the limiting procedure [9], we obtain the scaling dimension $\nu_k^\pm$ given by

$$6\left(\nu_k^\pm\right)^2 = \left(\sqrt{3}p \pm k\right)^2 - \frac{q^2}{2} \ , \tag{24}$$

to be compared with (5). The self energy $\Sigma(\omega, k)$ at the Fermi surface remains

$$\Sigma(\omega, k) \sim c\omega^{2\nu_k} \ . \tag{25}$$

For our system with fixed fermion charge and mass, ν_k and the fluid character is determined by the strength of the dipole interaction p.

To study explicit solutions, we set the fermion charge to $q = 1$, and numerically solve the flow equation (19) to determine where the system possesses a Fermi surface and the type of excitations. Figure 1 displays our results for $\nu_k^\pm$ evaluated at the Fermi momentum k_F. We recover the results of [15, 16] and classify the excitations as Fermi liquid ($p \lesssim -.53$), marginal Fermi liquid ($p \sim -.53$), non-Fermi liquid ($-.53 \lesssim p \lesssim .41$) and log-oscillatory ($p \gtrsim .41$). The system has a Fermi surface with $p \lesssim .41$ but no Fermi surface for larger values of p.

[1] Since the appearance of this work, the duality between poles and zeroes has subsequently been shown as a consequence of the two possible quantizations [45].



FIG. 2: Plots of $\Re \det G_R$ ($\Im \det G_R = 0$) with $q = 1$ for $p = -5$ (first panel) showing a pole at $k = k_F \approx 1.5$, and $p = 5$ (second panel) showing a zero at $k = k_L \approx 1.5$.

IV. DUALITY OF POLES AND ZEROES

Using the properties of the solutions (21) of the flow equation (19) we will show that there is a duality between poles and zeroes. We first emphasize that a pole of $G_\pm$ at $\omega = 0$ (eq. (23)) is not necessarily a pole of the determinant

$$\det G_R = G_+ G_- \ . \tag{26}$$

It is known [33] that in the conventional case, $p = 0$,

$$\det G_R(\omega = 0, k; p = 0) = 1 \ , \tag{27}$$

therefore it possesses neither poles nor zeroes. This is because poles (zeroes) of G_+ are cancelled by zeroes (poles) of G_- at the same momentum. We shall see that this coincidence of poles and zeroes is lifted when the dipole coupling is turned on, resulting in poles and zeroes of $\det G_R$.

It is easily deduced from the flow equation (19), that the reciprocal of $\xi_\pm$,

$$\zeta_\pm = \frac{1}{\xi_\pm} \ , \tag{28}$$

satisfies the flow equation

$$h\zeta'_\pm + \left[-\omega - \mu q(1-z) + \sqrt{h}\,(\pm k + \mu p z)\right]\zeta_\pm^2 - \mu q(1-z) + \sqrt{h}\,(\mp k - \mu p z) - \omega = 0 \ . \tag{29}$$

Upon comparison of the two flow equations, we see that $\zeta_\pm$ solves the same equation as $-\xi_\pm$ under the change of parameters $k \to -k$ and $p \to -p$. It follows that the inverse Green



FIG. 3: Plots of $\Re \det G_R$ (solid lines) and $\Im \det G_R$ (dashed lines) with $q = 1$ for $p = -0.1$ (first panel) showing a pole at $k = k_F \approx 0.8$ and a zero at $k = k_L \approx 1.0$, and $p = 0.1$ (second panel) showing a pole at $k = k_F \approx 1.0$ and a zero at $k = k_L \approx 0.8$.

function $G_{\pm}^{-1}(0, k)$ at p is identified with $-G_{\pm}(0, -k)$ at opposite dipole coupling $-p$. Using the relation between the two Green functions (22), we arrive at

$$\det G_R(\omega = 0, k; p) = \frac{1}{\det G_R(\omega = 0, k; -p)} \ . \tag{30}$$

This is the central result of our work and its importance relies on the fact that it establishes a relation between poles and zeroes of the determinant of the Green function at zero frequency between systems of opposite dipole coupling. For $p = 0$, we recover eq. (27). For $p \neq 0$, the poles found for $p < 0$ (Fermi liquid) correspond to zeroes for $p > 0$ (Mott insulator).

For large negative dipole coupling strength, i.e., $p \lesssim -.53$, we are in the Fermi liquid phase, since $\nu_k > 1/2$ (figure 1). In this regime, we obtain poles of $\det G_R$ and no zeroes. A typical example is shown in figure 2 (first panel) for $p = -5$. In this case, $\Im \det G_R$ vanishes and the graph has two poles at $k = k_F \approx \pm 1.5$, and no zeroes. According to the duality (30), we expect to see two zeroes at $k = k_L \approx \pm 1.5$ and no poles for a system with $p = 5$. This is indeed what we obtain by a numerical calculation of $\det G_R$ (second panel of figure 2). This is in the Mott insulator regime and no Fermi surface is found.

For small values of the dipole coupling strength ($|p| \lesssim .41$), we expect both zeroes and poles. Again, using the duality (30), the zeroes can be deduced from the poles at opposite p (and vice versa). An example is shown in figure 3 for $p = \pm .1$. Unlike in the case of large p, $\Im \det G_R$ does not vanish. Instead, the zeroes for $p = .1$ found at $k = k_L \approx \pm .8$ are isolated zeroes of both $\Re \det G_R$ and $\Im \det G_R$. Correspondingly, the same is true for the poles at $k = k_F \approx \pm .8$ for $p = -.1$. At the other set of zeroes at $k = k_L \approx \pm 1.0$ for $p = -.1$ (and correspondingly poles at $k = k_F \approx \pm 1.0$ for $p = .1$), $\Im \det G_R$ vanishes over a range



FIG. 4: Poles at $k = k_F$ (blue lines) and zeroes at $k = k_L$ (red lines) *vs.* p with $q = 1$. Notice the symmetry under $k \to -k$, and the duality of poles and zeroes under $p \to -p$.

containing the zeroes (poles).

Finally, figure 4 shows the location of poles ($k = k_F$) and zeroes ($k = k_L$) as the dipole coupling strength p varies. Notice the symmetry under $k \to -k$, as well as the interchange $k_L \leftrightarrow k_F$ under the mapping $p \to -p$.

V. CONCLUSIONS AND DISCUSSION

We have shown that a holographic theory with a bulk dipole interaction between a massless fermion and gauge field possesses a robust phase diagram including Fermi and non-Fermi liquids, insulating Mott state and pseudo-gap state. The various phases are identified by the structure of the poles and zeroes found in G_R. The selection of the phase is controlled by the strength of the dipole coupling which plays the role of an order parameter in the holographic system [15, 16].

A pole of G_R is indicative of a Fermi or non-Fermi fluid while a zero is responsible for an insulating phase. It is the coexistence of both that underlies the identification of a holographic pseudo-gap state. We showed that a duality exists relating systems of opposite dipole coupling strength p. This duality maps zeroes to poles and vice versa, pointing to the interesting possibility of understanding the properties of a system with zeroes (insulating phase) by mapping the system to one with poles (Fermi liquid).

It will be interesting to explore further the pseudo-gap state and other aspects that may be recovered by the evolution of the Fermi and Luttinger surfaces. Of particular significance will be the response of the system with $\vec{k}$ dependence. Also, it might be worth exploring

further the physical significance of the behavior of $\Im \det G_R$ near zeroes and poles.

As we discussed in the introduction, the interplay between the state of matter and the appearance of poles and zeroes is also observed in QCD. The difference with QCD is that the appearance of free quarks or bound states depends on the dynamics of the strong gauge theory while in our case there is a variation of a simple coupling that gives the various phases of matter. It would be intriguing to find other dynamical systems exhibiting this behavior.

One may wonder if this analysis can be applied to holographic superconductors. In BCS superconductivity the role of the order parameter is played by the condensate of a scalar field. The formation of a condensate corresponds to a critical temperature below which the system enters the superconductivity phase. In this phase, the electrons are combined to form Cooper pairs which are electronic bound states and therefore correspond to zeroes. On the contrary, the electrons in the normal phase are free corresponding to poles.

In spite of the similarities with superconductivity, our analysis does not explicitly break a gauge symmetry that could have defined an order parameter for the various phases. Instead the metallic phases are controlled by an explicit coupling of charged fermions to a gauge field. Nevertheless, it will be enlightening to consider how a bulk scalar responsible for superconductivity or the coupling will influence the system. Also worth exploring is if a description with additional fields could explain the Green function zeroes in terms of a composite excitation.

Another way to better understand our results is to calculate the holographic entanglement entropy in our theory. The entanglement entropy [46, 47] has proven to be a powerful tool in counting the degrees of freedom available in a holographic system. In [48] it was found that the holographic entanglement entropy of a superconducting phase is less than that of the normal phase due to the formation of Cooper pairs reducing the degrees of freedom available (see also [49]). Near the contact interface, the normal phase entanglement entropy possesses a higher value compared to the superconducting phase due to the proximity effect: the leakage of Cooper pairs to the normal phase so more free electrons were available. Similarly we expect that the entanglement entropy in a holographic fermionic system with a dipole coupling will vary, as the degrees of freedom change from the Fermi liquid phase to the Mott insulating phase providing important information on the various phases of the system. However, such a study has to overcome the basic difficulty of working with a probe fermonic system. The boundary theory will see the effect of the dipole coupling only with a fully

back-reacted solution. Once a full solution is obtained the dipole coupling could leave some trace on the metric. Work in this direction is in progress.

Acknowledgments

We thank P. W. Phillips for illuminating discussions. E.P is partially supported by the Greek Ministry of Education and Religious Affairs, Sport and Culture through the ARISTEIA II action of the operational program Education and Lifelong Learning.

---

[1] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. **2** (1998) 231 [Int. J. Theor. Phys. **38** (1999) 1113] [arXiv:hep-th/9711200].

[2] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, "Building a Holographic Superconductor," Phys. Rev. Lett. **101**, 031601 (2008) [arXiv:0803.3295]; S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, "Holographic Superconductors," JHEP **0812**, 015 (2008) [arXiv:0810.1563].

[3] M. Cubrovic, J. Zaanen and K. Schalm, "String Theory, Quantum Phase Transitions and the Emergent Fermi-Liquid," Science **325**, 439 (2009). [arXiv:0904.1993 [hep-th]].

[4] S. Bhattacharyya *et al*, V. E. Hubeny, S. Minwalla and M. Rangamani, "Nonlinear Fluid Dynamics from Gravity," JHEP **0802**, 045 (2008). [arXiv:0712.2456 [hep-th]].

[5] N. Iqbal, H. Liu, M. Mezei and Q. Si, "Quantum phase transitions in holographic models of magnetism and superconductors," Phys. Rev. D **82**, 045002 (2010). [arXiv:1003.0010 [hep-th]].

[6] C. P. Herzog, P. Kovtun, S. Sachdev and D. T. Son, "Quantum critical transport, duality, and M-theory," Phys. Rev. D **75**, 085020 (2007). [arXiv:hep-th/0701036].

[7] G. Koutsoumbas, E. Papantonopoulos and G. Siopsis, "Exact Gravity Dual of a Gapless Superconductor," JHEP **0907**, 026 (2009), [arXiv:0902.0733 [hep-th]]; Q. Pan, B. Wang, E. Papantonopoulos, J. Oliveira and A. B. Pavan, "Holographic Superconductors with various condensates in Einstein-Gauss-Bonnet gravity," Phys. Rev. D **81**, 106007 (2010). [arXiv:0912.2475 [hep-th]]; G. Siopsis and J. Therrien, "Analytic Calculation of Properties of Holographic Superconductors," JHEP **1005**, 013 (2010), [arXiv:1003.4275 [hep-th]].

[8] H. Liu, J. McGreevy and D. Vegh, "Non-Fermi liquids from holography," Phys. Rev. D **83**, 065029 (2011) [arXiv:0903.2477 [hep-th]].

[9] T. Faulkner, H. Liu, J. McGreevy and D. Vegh, "Emergent quantum criticality, Fermi surfaces, and AdS(2)," Phys. Rev. D **83**, 125002 (2011) [arXiv:0907.2694 [hep-th]].

[10] S. A. Hartnoll, J. Polchinski, E. Silverstein and D. Tong, "Towards strange metallic holography," JHEP **1004**, 120 (2010) [arXiv:0912.1061 [hep-th]].

[11] S. A. Hartnoll and A. Tavanfar, "Electron stars for holographic metallic criticality," Phys. Rev. D **83**, 046003 (2011) [arXiv:1008.2828 [hep-th]].

[12] A. Allais and J. McGreevy, "How to construct a gravitating quantum electron star," Phys. Rev. D **88**, no. 6, 066006 (2013) [arXiv:1306.6075 [hep-th]].

[13] M. Cubrovic, J. Zaanen and K. Schalm, "Constructing the AdS Dual of a Fermi Liquid: AdS Black Holes with Dirac Hair," JHEP **1110**, 017 (2011) [arXiv:1012.5681 [hep-th]].

[14] S. Sachdev, "A model of a Fermi liquid using gauge-gravity duality," Phys. Rev. D **84**, 066009 (2011) [arXiv:1107.5321 [hep-th]].

[15] M. Edalati, R. G. Leigh and P. W. Phillips, "Dynamically Generated Mott Gap from Holography," Phys. Rev. Lett. **106**, 091602 (2011) [arXiv:1010.3238 [hep-th]].

[16] M. Edalati, R. G. Leigh, K. W. Lo and P. W. Phillips, "Dynamical Gap and Cuprate-like Physics from Holography," Phys. Rev. D **83**, 046012 (2011) [arXiv:1012.3751 [hep-th]].

[17] W. J. Li and H. Zhang, "Holographic non-relativistic fermionic fixed point and bulk dipole coupling," JHEP **1111**, 018 (2011) [arXiv:1110.4559 [hep-th]].

[18] X. M. Kuang, B. Wang and J. P. Wu, "Dipole Coupling Effect of Holographic Fermion in the Background of Charged Gauss-Bonnet AdS Black Hole," JHEP **1207**, 125 (2012) [arXiv:1205.6674 [hep-th]].

[19] L. Q. Fang, X. H. Ge and X. M. Kuang, "Holographic fermions with running chemical potential and dipole coupling," Nucl. Phys. B **877**, 807 (2013) [arXiv:1304.7431 [hep-th]].

[20] J. P. Wu, "The charged Lifshitz black brane geometry and the bulk dipole coupling," Phys. Lett. B **728**, 450 (2014).

[21] P. W. Phillips, B. W. Langley, J. A. Hutasoit, "Un-Fermi liquids: Unparticles in strongly correlated electron matter," Phys. Rev. B **88**, 115129 (2013), [arXiv:1305.0006 [cond-mat]].

[22] D. Guarrera and J. McGreevy, "Holographic Fermi surfaces and bulk dipole couplings," [arXiv:1102.3908 [hep-th]].

[23] J. M. Luttinger, "Fermi Surface and Some Simple Equilibrium Properties of a System of Interacting Fermions," Phys. Rev. **119**, 1153 (1960).

[24] T. D. Stanescu, P. W. Phillips, and T. Choy, P "Theory of the Luttinger surface in doped Mott insulators," hys. Rev. B **75**, 104503 (2007), [arXiv:cond-mat/0602280].

[25] I. Dzyaloshinskii, "Some consequences of the Luttinger theorem: The Luttinger surfaces in non-Fermi liquids and Mott insulators," Phys. Rev. B **68**, 085113 (2003).

[26] G. 't Hooft, "A Two-Dimensional Model for Mesons," Nucl. Phys. B **75**, 461 (1974).

[27] T. D. Stanescu and G. Kotliar, "Fermi arcs and hidden zeros of the Green function in the pseudogap state," Phys. Rev. B **74**, 125110 (2006), [arXiv:cond-mat/0508302]; R. M. Konik, T. M. Rice, and A. M. Tsvelik, "Doped Spin Liquid: Luttinger Sum Rule and Low Temperature Order," Phys. Rev. Lett. **96**, 086407 (2006), [arXiv:cond-mat/0511268].

[28] S. Hong and P. W. Phillips, "Towards the standard model for Fermi arcs from a Wilsonian reduction of the Hubbard model," Phys. Rev. B **86**, 115118 (2012), [arXiv:1110.0440 [cond-mat.str-el]].

[29] S. Sakai, Y. Motome, and M. Imada, "Evolution of Electronic Structure of Doped Mott Insulators: Reconstruction of Poles and Zeros of Green?s Function," Phys. Rev. Lett. **102**, 056404 (2009), [arXiv:0809.0950v1 [cond-mat.str-el]].

[30] S. Sakai, Y. Motome, and M. Imada, "Doped high-Tc cuprate superconductors elucidated in the light of zeros and poles of the electronic Green?s function," Phys. Rev. B **82**, 134505 (2010), [arXiv:1004.2569 [cond-mat.str-el]].

[31] K. B. Dave, P. W. Phillips and C. L. Kane, "Absence of Luttinger's Theorem," Phys. Rev. Lett. **110**, 090403 (2013) [arXiv:1207.4201 [cond-mat.str-el]].

[32] S. S. Lee, "A Non-Fermi Liquid from a Charged Black Hole: A Critical Fermi Ball," Phys. Rev. D **79**, 086006 (2009) [arXiv:0809.3402 [hep-th]].

[33] T. Faulkner, N. Iqbal, H. Liu, J. McGreevy and D. Vegh, "From Black Holes to Strange Metals," [arXiv:1003.1728 [hep-th]].

[34] S. S. Gubser, F. D. Rocha and A. Yarom, "Fermion correlators in non-abelian holographic superconductors," JHEP **1011**, 085 (2010) [arXiv:1002.4416 [hep-th]]; M. Ammon, J. Erdmenger, M. Kaminski and A. O'Bannon, "Fermionic Operator Mixing in Holographic p-wave Superfluids," JHEP **1005**, 053 (2010) [arXiv:1003.1134 [hep-th]].

[35] F. Benini, C. P. Herzog and A. Yarom, "Holographic Fermi arcs and a d-wave gap," Phys. Lett. B **701**, 626 (2011) [arXiv:1006.0731 [hep-th]].

[36] Y. Liu, K. Schalm, Y. W. Sun and J. Zaanen, "Lattice Potentials and Fermions in Holographic non Fermi-Liquids: Hybridizing Local Quantum Criticality," JHEP **1210**, 036 (2012) [arXiv:1205.5227 [hep-th]].

[37] S. S. Gubser, C. P. Herzog, S. S. Pufu and T. Tesileanu, "Superconductors from Superstrings," Phys. Rev. Lett. **103**, 141601 (2009) [arXiv:0907.3510 [hep-th]].

[38] J. P. Gauntlett, S. Kim, O. Varela and D. Waldram, "Consistent supersymmetric Kaluza–Klein truncations with massive modes," JHEP **0904** (2009) 102 [arXiv:0901.0676 [hep-th]].

[39] J. P. Gauntlett, J. Sonner and T. Wiseman, "Holographic superconductivity in M-Theory," Phys. Rev. Lett. **103** (2009) 151601 [arXiv:0907.3796 [hep-th]].

[40] J. P. Gauntlett, J. Sonner and T. Wiseman, "Quantum Criticality and Holographic Superconductors in M-theory," JHEP **1002**, 060 (2010) [arXiv:0912.0512 [hep-th]].

[41] N. Bobev, N. Halmagyi, K. Pilch and N. P. Warner, "Supergravity Instabilities of Non-Supersymmetric Quantum Critical Points," Class. Quant. Grav. **27**, 235013 (2010); N. Bobev, A. Kundu, K. Pilch and N. P. Warner, "Minimal Holographic Superconductors from Maximal Supergravity," JHEP **1203**, 064 (2012).

[42] I. Bah, A. Faraggi, J. I. Jottar, R. G. Leigh and L. A. Pando Zayas, "Fermions and $D = 11$ Supergravity On Squashed Sasaki-Einstein Manifolds," JHEP **1102**, 068 (2011) [arXiv:1008.1423 [hep-th]].

[43] I. Bah, A. Faraggi, J. I. Jottar and R. G. Leigh, "Fermions and Type IIB Supergravity On Squashed Sasaki-Einstein Manifolds," JHEP **1101**, 100 (2011) [arXiv:1009.1615 [hep-th]].

[44] O. DeWolfe, S. S. Gubser, and C. Rosen, "Fermi surfaces in N=4 Super-Yang-Mills theory," Phys. Rev. D **86**, (2012) 106002 , [arXiv:1207.3352 [hep-th]]; O. DeWolfe, S. S. Gubser, and C. Rosen,, "Minding the Gap in N=4 Super-Yang-Mills," [arXiv:1312.7347 [hep-th]].

[45] G. Vanacore, P. W. Phillips, "Minding the Gap in Holographic Models of Interacting Fermions," Phys. Rev. D **90**, 044022 (2014), [arXiv:1405.1041 [cond-mat]].

[46] S. Ryu and T. Takayanagi, "Holographic derivation of entanglement entropy from AdS/CFT," Phys. Rev. Lett. **96**, 181602 (2006) [hep-th/0603001].

[47] S. Ryu and T. Takayanagi, "Aspects of Holographic Entanglement Entropy," JHEP **0608**, 045 (2006) [hep-th/0605073].

[48] X. M. Kuang, E. Papantonopoulos and B. Wang, "Entanglement Entropy as a Probe of the Proximity Effect in Holographic Superconductors," JHEP **1405**, 130 (2014) [arXiv:1401.5720 [hep-th]].

[49] T. Albash and C. V. Johnson, "Holographic Studies of Entanglement Entropy in Superconductors," JHEP **1205**, 079 (2012) [arXiv:1202.2605 [hep-th]].